# Adolescent Health in Arab countries: Statistics and Social context

Doha, 3 December 2013

Carla Makhlouf Obermeyer Director, Center for Research on Population and Health Faculty of Health Sciences, American University of Beirut

### **Outline**


- 1. General demographic and epidemiologic evidence
- 2. Main health risks among adolescents
- 3. The social context of adolescent health

## Part 1. Demographic and epidemiological evidence

## 1.1. More adolescents, longer adolescence

- 1. "Youth bulge:" >1/2 population under 25
- 2. Marriage: later and less frequent
- → Expansion of the adolescence transition





Sources: National surveys between 2003-2009 (Loaiza and Wong 2012; PRB 2013

### 1.2. Burden of disease: causes of death (over 5%)


#### Top causes:

Cardiovascular
Transport
Unintentional injuries
Diarrhea/ infections
Cancer, diabetes

globalglobal

about equal

- < global
- > global



Source: Institute of Health Metrics and Evaluation: Global Burden of Disease study 2013


### 1.3. Burden of disease: causes of DALYs (over 5%)

Mental/ behavioral disorders, 25% DALYs Musculo-skeletal, nutritional, transport, respiratory, cardiovascular and diabetes Diarrhea/ infections

> global

all > global


< global



Source: Institute of Health Metrics and Evaluation: Global Burden of Disease study 2013

## Part 2. Main health risks among adolescents


### 2.1. Overweight and obesity are high



The nutrition transition takes its toll Countries at different stages



## 2.2. Physical activity is low



Global School Health Survey, Guthold et al 2010

- Insufficient activity, many hours sitting
- Especially women





## 2.3. Transport injuries: much worse than elsewhere

# Far worse in Arab countries than rest of the world

2<sup>nd</sup> leading cause of death among males 2<sup>nd</sup> highest mortality rate (LMICs) Highest among males 15-29 Trauma data confirms


Bad driving, speed, violating traffic laws Low seat belt use







## 2.4. Smoking: too high, especially among males










## 2.5. Mental health, links to violence?



1/4 DALYs

Surveys: anxiety, depression, sadness, hopelessness,

Exposure to personal violence School fighting: EMRO high %

Socially acceptable violence: honor crimes, wife beating Regional violence and war Adolescents vulnerable But critique of medicalization of violence

### 3. The social context of adolescent health

## Mismatches, disparities

### 3.1. Mismatch between education and employment

- Progress in education but not jobs
- Uncompetitive market, nepotism and corruption
- Highest youth employment in the world, at all levels
- Alienation, despair, migration?

### 3.2. Gender: improvement and disparities

- Progress in women's education, greater opportunities
- But worst Global Gender Gap indicators: lack of employment and no participation in politics
- Legal reforms, but still limited legal protection for women

### 3.3. Reproductive health

- Delayed marriage, premarital sex
- Insufficient information and protection

## Aspirations, confrontation, convergence

- Political movements: participation of youth
- Difficult choices: family/ emigration
- Democracy / fundamentalism
- The political generation gap
- Social media

#### Youth who think they will move away (%)

